

jingcheng zhang      ORCID iD: 0000-0002-7555-8848

## Fecal specimen diagnosis 2019 Novel Coronavirus–Infected Pneumonia

JingCheng Zhang, SaiBin Wang, YaDong Xue

Department of respiration, Jinhua Hospital of Zhejiang University(Jinhua  
Municipal Central Hospital), Jinhua, China 321000

### **Abstract**

The emergence and spread of 2019 Novel Coronavirus–Infected Pneumonia (COVID-19) from Wuhan, China, it has spread globally. We extracted the data on 14 patients with laboratory-confirmed COVID-19 from Jinhua Municipal Central hospital through January 27th, 2020. We found that compared to pharyngeal swab specimens, nucleic acid detection of COVID-19 in fecal specimens was equally accurate. And we found that patients with a positive stool test did not experience gastrointestinal symptoms and had nothing to do with the severity of the lung infection. These results may help to understand the clinical diagnosis and the changes of clinical parameters of COVID-19.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1002/jmv.25742.

This article is protected by copyright. All rights reserved.

## 1 INTRODUCTION

In December 2019, a new coronavirus (COVID-19) pneumonia outbreak occurred in Wuhan, China<sup>1</sup>. Since the virus was detected in December 2019, more than 77,000 people have been infected in China, causing more than 2,600 deaths<sup>2</sup>. Globally, cases have been reported in 33 countries and 5 continents<sup>3</sup>. Patients' clinical manifestations included fever, cough, dyspnea, myalgia, fatigue, and chest computed tomography were ground-glass opacity<sup>4</sup>. A novel coronavirus currently named 2019 novel coronavirus (COVID-19), RT-PCR method can be used for the detection of COVID-19 in oral swabs<sup>5</sup>. In SARS-CoV and MERS-CoV infected patients, intestinal infections are very common<sup>6,7</sup>. Results have now confirmed the presence of live virus in stool samples from patients with COVID-19<sup>8</sup>, which means that stool samples may contaminate hands, food, and water, and may cause infection by invading the mouth, respiratory mucosa, and conjunctiva. We report detection method of 14 cases with confirmed COVID-19 infection admitted to hospitals in Jinhua, aiming to investigate the other possible transmission route of this virus.

## 2 METHODS

We performed a retrospective analysis of 14 laboratory-confirmed cases with COVID-19 pneumonia. Data were obtained from Department of Infectious Diseases Jinhua Hospital of Zhejiang University. Patients were hospitalized from January 27, 2020, to February 10, 2020, with final follow-up for this report on February 9, 2020. Early we are confirmed cases with COVID-19 pneumonia were a real-time reverse-transcriptase polymerase-chain-reaction (RT-PCR) assay for


pharyngeal swab specimens, and that all oropharyngeal swabs specimens were collected by a senior infectious physician with more than 10 years of experience. Starting from February 4, 2020, we have added stool samples to all cases.

### 3 RESULTS

The median age of the patients was 41 years ( 18-87 years), and 7 (50%) were female. The most common symptoms include fever (92.8%) and cough (71.4%), no fever is a 87-year-old female patient, all patients had no vomiting and diarrhea. 4 patients(28.5%) returned from Wuhan, the remaining 10 (71.4%) had a history of contact with COVID-19 pneumonia patients.

Since February 4th, we have collected patient samples intensively as the supply of PCR kits has increased. It was found that 5 of the 14 confirmed patients (35.7%) had a positive stool sample COVID-19 nucleic acid, and patients with positive stool samples were also positive for oropharyngeal swabs specimens at least the day before (Figure 1); The same trend is that patients with negative stool samples are also negative for oropharyngeal swabs for at least the first two days(Figure 1). This may indicate that the accuracy of fecal specimen detection and oropharyngeal swabs is the same.

Figure 1. Viral nucleic acid test results in 14 patients with COVID-19.



All 14 patients had computed tomography on admission, 92.8% manifested as pneumonia. 1 patient (Patient 3) chest computed tomography images did not demonstrate any consolidation or scarring (Figure 2), however, this patient was positive for multiple stool and oropharyngeal swabs specimens (Figure 1). Among the remaining 13 patients, ground glass opacity was observed in both lungs. 2 patients (Patient 6 and Patient 14) severely yielded more prominent abnormalities on chest computed tomography than non-severe cases (Figure 3), the nucleic acid test was negative for stool samples in these two patients, this indicates that fecal specimens are not related to nucleic acid positivity and severity. As of February 9, 2020, 3 patients recovered, but 11 were still being treated in the hospital.

Figure 2. Chest computed tomographic images, February 5, 2020 (Illness Day 10, Hospital Day 6).





Figure 3. Chest computed tomographic images. Shown Patient are chest images obtained at on Day 5 ( A and B), and another patient Day 5 ( C and D) after admission.



#### **4 DISCUSSION**

According to latest pilot experiment, 4 out of 62 stool specimens (6.5%) tested positive to COVID-19<sup>9</sup>, but we found very high accuracy in detecting nucleic acids in stool samples. Recently, reports of fecal nucleic acid tests to diagnose COVID-19 have increased<sup>10,11</sup>. In our cases, we found that patients with a positive stool test did not experience gastrointestinal symptoms and had nothing to do with the severity of the lung infection. Stool specimen collection will reduce medical staff infections compared to oropharyngeal swabs specimens, especially when protective equipment in China is inadequate. We are not sure if there is fecal-oral transmission, but fomite transmission might have contributed considerably to the rapid spread of COVID-19. Finally, this report highlights the need of health protection, which should be considered for transmission through stool, these discoveries will curb the rapid spread of COVID-19 worldwide.

#### **Conflicts of interest**

The author has no conflicts of interest to disclose.

#### **Acknowledgement**

The author thanks Dr. Saibin Wang for sharing her personal observations.

## References

1. Zhu N, Zhang D, Wang W, et al. A novel coronavirus from patients with pneumonia in China. *N Engl J Med*.2020.
- 2.National Health Commission of the People’s Republic of China. Assessed on February 25th, 2020.(<http://www.nhc.gov.cn/xcs/yqfkdt/202002/945bd98a9d884aeeb54d76afa02ca813.shtml>)
3. World Health Organization. Novel coronavirus(2019-nCoV): situation report-36. Accessed February 25, 2020. ([https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200225-sitrep-36-covid-19.pdf?sfvrsn=2791b4e0\\_2](https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200225-sitrep-36-covid-19.pdf?sfvrsn=2791b4e0_2))
- 4.Wang D, Hu B, Hu C, et al. Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus–Infected Pneumonia in Wuhan, China. *JAMA*. 2020. doi: 10.1001/jama.2020.1585.
5. Zhou P, Yang XL, Wang XG, et al. Discovery of a novel coronavirus associated with the recent pneumonia outbreak in humans and its potential bat origin. *Nature*.2020. doi: 10.1038/s41586-020-2012-7.
6. Shi X, Gong E, Gao D, et al. Severe acute respiratory syndrome associated coronavirus is detected in intestinal tissues of fatal cases. *Am J Gastroenterol*. 2005;100(1):169–176.
7. Ding Y, He L, Zhang Q, et al. Organ distribution of severe acute respiratory syndrome (SARS) associated coronavirus (SARS-CoV) in SARS patients: implications for pathogenesis and virus transmission pathways. *J Pathol*. 2004;203(2):622–630.
8. Zhang Y, Chen H, Zhu S, et al. Isolation of 2019-nCoV from a Stool Specimen of a Laboratory-Confirmed Case of the Coronavirus Disease 2019 (COVID-19). *China CDC Weekly*. 2020, 2(8): 123-124.

9. Guan W, Ni Z, Hu Y, et al. Clinical characteristics of 2019 novel coronavirus infection in China. Medrxiv. 2020.
10. Zhang H, Kang Z, Gong H, et al. The digestive system is a potential route of 2019-nCov infection: a bioinformatics analysis based on single-cell transcriptomes. bioRxiv .2020.
11. Zhang W, Du R, Li B, Bing Yan, et al. Molecular and serological investigation of 2019-nCoV infected patients: implication of multiple shedding routes. Emerging Microbes Infect. 9: 386-389;2020.